



Reactive oxygen species target specific tryptophan site in the mitochondrial ATP synthase

Sascha Rexroth^{a,*}, Ansgar Poetsch^a, Matthias Rögner^a, Andrea Hamann^b, Alexandra Werner^b, Heinz D. Osiewacz^b, Eva R. Schäfer^c, Holger Seelert^c, Norbert A. Dencher^c

^a Plant Biochemistry, Faculty of Biology & Biotechnology, Ruhr University Bochum, D-44780 Bochum, Germany

^b Faculty for Biosciences, Molecular Developmental Biology, Cluster of Excellence Macromolecular Complexes, Johann Wolfgang Goethe University, Max-von-Laue-Str. 9, 60438 Frankfurt, Germany

^c Physical Biochemistry, Department of Chemistry, Technische Universität Darmstadt, Petersenstrasse 22, D-64287 Darmstadt, Germany

ARTICLE INFO

Article history:

Received 12 September 2011

Received in revised form 4 November 2011

Accepted 10 November 2011

Available online xxxx

Keywords:

ATP synthase

Oxidative stress

Podospira anserina

Ageing

Posttranslational modification

ABSTRACT

The release of reactive oxygen species (ROS) as side products of aerobic metabolism in the mitochondria is an unavoidable consequence. As the capacity of organisms to deal with this exposure declines with age, accumulation of molecular damage caused by ROS has been defined as one of the central events during the ageing process in biological systems as well as in numerous diseases such as Alzheimer's and Parkinson's Dementia. In the filamentous fungus *Podospira anserina*, an ageing model with a clear defined mitochondrial etiology of ageing, in addition to the mitochondrial aconitase the ATP synthase alpha subunit was defined recently as a hot spot for oxidative modifications induced by ROS. In this report we show, that this reactivity is not randomly distributed over the ATP Synthase, but is channeled to a single tryptophan residue 503. This residue serves as an intra-molecular quencher for oxidative species and might also be involved in the metabolic perception of oxidative stress or regulation of enzyme activity. A putative metal binding site in the proximity of this tryptophan residue appears to be crucial for the molecular mechanism for the selective targeting of oxidative damage.

© 2011 Published by Elsevier B.V.

1. Introduction

Molecular damage caused by the exposure to ROS is a major cause of numerous diseases such as Alzheimer's and Parkinson's Dementia as well as a central factor of the ageing process by the 'free radical theory of ageing' (FRTA) [1] and its refined version, the 'mitochondrial free radical theory of ageing' (MFRTA) [2]. The latter theory identifies mitochondria as the major site of ROS generation and the predominant target of damage of all kinds of biomolecules. As a result, mitochondria become dysfunctional and biological systems degenerate and die. On the other hand, however, ROS are also employed in many life-sustaining mechanisms such as in signaling pathways and defense against pathogens.

Abbreviations: Desferal, Deferoxamine N'-[5-[acetyl(hydroxyamino)pentyl]-N-[4-[(5-aminopentyl)(hydroxyamino)-4-oxobutanoyl]amino]pentyl]-N-hydroxysuccinamide; FRTA, Free radical theory of ageing; Kyn, kynurenine; MFRTA, mitochondrial free radical theory of ageing; MS, mass spectrometry; NFK, N-formyl-kynurenine; ROS, reactive oxygen species; SRM, single reaction monitoring; TRP-OH, hydroxy-tryptophan.

* Corresponding author at: Plant Biochemistry, Faculty of Biology & Biotechnology, Ruhr University Bochum, D-44780 Bochum, Germany. Tel.: +49 0234 32 29896; fax: +49 0234 32 14322.

E-mail address: sascha.rexroth@rub.de (S. Rexroth).

In ageing research the conditions and effects of ROS generation, signaling, scavenging and damaging, are intensively studied in various systems including different model organisms. *Podospira anserina* is one of these organisms. This filamentous fungus is characterized by a short lifespan of a few weeks and is tractable to experimentation. It contains a small genome [3], of which the sequence has been completely determined [4]. Ageing of *P. anserina* has a clear mitochondrial etiology. A number of specific factors and pathways which are basically involved in keeping a functional population of mitochondria have been demonstrated to affect ageing and lifespan [3, 5–10]. Although ROS generation and scavenging do clearly contribute to the ageing process in *P. anserina* [11, 12], recently counter-intuitive data were obtained when the effect of the modulation of mitochondrial superoxide dismutase (PaSOD3) was studied [13]. In contrast to what is expected from the MFRTA, strains deleted for *PaSod3* were not short-lived and strains overexpressing *PaSod3* were not long- but short-lived. Intriguingly, the latter strains were characterized by a reduced abundance of a mitochondrial peroxiredoxin involved in scavenging of hydrogen peroxide, and in PaCLP protease, as a part of the mitochondrial protein quality control system. Taking these observations into account allowed the generation of a mathematical model that, at least partly, explains the unexpected results obtained from the study (manuscript in preparation). In particular, in addition to ROS metabolism mitochondrial protein quality control

systems need to be included in networks of pathways involved in the control of ageing. Such systems may be triggered by and acting on specific posttranslational modifications. One of such modifications has also been identified in *P. anserina* mitochondria as the irreversible oxidation of tryptophan residues and the formation of *N*-formylkynurenine (NFK) [14, 15]. NFK is frequently observed in proteins by mass spectrometry along with the other oxidation intermediates displayed in Fig. 1, particularly in tissues with high metabolic rates and in proteins with long half lives [16]. Prominent is its presence in human and bovine heart mitochondrial proteins [15, 17], rat skeletal muscle proteins [18], bovine α -crystalline [19], as well as CP43 [20] and LHClI [21] in photosynthetic membranes.

Here we show that oxidative modifications are not—as intuitively accepted—randomly distributed to all tryptophan residues over the entire sequence of the alpha subunit of the mitochondrial ATPase, but selectively targeted to one specific tryptophan.

2. Materials and methods

2.1. *P. anserina* cultivation and isolation of crude mitochondria fraction

P. anserina is a filamentous ascomycete which is normally growing on solid medium. To obtain sufficient material for mitochondria isolation, after germination of monokaryotic ascospores on solid cornmeal medium with 60 mM ammonium acetate, small pieces of mycelium are first grown 2–3 days on solid PASM medium [22] under permanent light. To prevent transfer of solid medium into the subsequently used liquid medium, the solid medium was overlaid with a cellophane sheet. The mycelium from this solid medium was then transferred to liquid CM medium [23] and incubated for 2 days shaking under permanent light. Different age stages were obtained by pre-growth to the respective age on solid medium and subsequent culture in liquid medium. For instance, to isolate mitochondria from 6 days old cultures, ascospores were germinated for two days on solid cornmeal medium with ammonium acetate. Subsequently, pieces of mycelium from the growth front were transferred to solid medium with cellophane. After two days of growth, the mycelium was scratched from the plates and used to inoculate liquid medium. After additional 2 days of growth the mycelium was harvested, giving a total age of an individual of 6 days. Senescent cultures were isolated by growing the fungus on solid medium until it reaches senescence (= stop of growth, alteration of pigmentation). The solid medium with cellophane was now inoculated with mycelium 2–3 cm (4–5 days) behind the growth front. To prevent rejuvenation, cultivation of senescent mycelium in liquid medium was performed without shaking in Fernbach flasks (permanent light, 5–7 days). The isolation of mitochondria from these different mycelial samples was modified according to Gredilla [12]: After high-speed centrifugation (15,000 g, 20 min) the mitochondria were directly resuspended in isolation buffer without BSA, additional wash steps were omitted. In total crude mitochondrial extracts of three independent wild-type

isolates of the age stages 6 days and senescent (14–15 days) were prepared. In control experiments 100 μ M Desferal was added to all buffers used during isolation.

2.2. 2-D Blue-native/tricine-SDS-PAGE and colorless-native PAGE

Isolated crude mitochondrial fractions were resuspended in a medium containing 50 mM imidazol, pH 7.0, 50 mM NaCl, 5 mM ϵ -aminocaproic acid, and 10% glycerol. The membranes were solubilized on ice for 30 min with the non-ionic detergent digitonin at a final concentration of 1% (4 g digitonin/g protein). Insoluble material was removed by centrifugation (21,000 g, 4 $^{\circ}$ C, 10 min). The supernatant was loaded directly onto blue-native gradient gels. Blue-native PAGE was performed using the Hoefer SE 600 system (18 \times 16 \times 0.15 cm³, 10 lanes) as described previously [24, 25]. Stacking gels with a total acrylamide concentration of 3.5% and separating gels with linear acrylamide gradients 5–13% were prepared. 250 μ g solubilized protein (as determined before solubilization) was applied per lane. Bovine heart mitochondria prepared from tissue and stored at -80° C (4 g digitonin/g protein, 70 μ g of protein before solubilization) served as molecular mass standard.

After electrophoresis, gels were stained with Coomassie R-250 or lanes of the BN-gel were cut out and incubated in a solution of 1% (w/v) SDS and 1% (w/v) β -mercaptoethanol at 20 $^{\circ}$ C for 30 min. Subsequently, lanes were analyzed by Tricine-SDS-PAGE in second dimension on a gel with two stacking gels, one native and one denaturing, with a total acrylamide concentration of 10% and a separating gel with 16.5% [26].

2.3. Protein analysis

After electrophoresis, gels were silver stained [27, 28]. Protein bands of interest were excised inside a laminar flow hood. Silver stained spots were destained [29], and in-gel digestion was performed with trypsin [30]. After extraction of the peptides with 50% (v/v) acetonitrile / 0.5% (v/v) formic acid the solvent was removed completely by lyophilization. Previous to LC-ESI-MS/MS analysis peptides were resuspended in 2% acetonitrile with 0.1% formic acid.

2.4. LC-MS analysis

LC-MS analysis was performed as described previously [31] with slight modifications. For reverse phase chromatography, a gradient of solvent A (95% water, 5% acetonitrile, 0.1% formic acid) and solvent B (10% water, 85% acetonitrile, 5% isopropanol, 0.1% formic acid) was used. For MS analysis, a Thermo LTQ Orbitrap mass spectrometer was operated in a duty cycle consisting of one 400–2000 *m/z* FT-MS and four MS/MS LTQ scans. For detection of low quantities of tryptophan oxidation for all three tryptophan sites in the ATP synthase alpha subunit an inclusion list was defined for the doubly and triply charged ions of the peptides ILQWEADFLSHLK, YSIVVAATASEAAPLQYLAPFTGASIGEWFR, and WNSGNDETK and their respective variants with oxidized tryptophan residues (Table 1).

2.5. Data analysis

Analysis of the LC-MS/MS data was performed using the Sequest algorithm [32] implemented in the Bioworks 3.3.1 software (Thermo scientific) for peptide identification versus a database [4] consisting of all *P. anserina* proteins with a tolerance of 10 ppm for the precursor mass accuracy and 1 u for the fragment mass accuracy. For detection of oxidized peptides for tryptophan residues modifications of 15.9949, 31.9898, and 3.9949 u were defined as parameters during the search for hydroxy-tryptophan, *N*-formyl kynurenin and kynurenin, respectively. False discovery rates were estimated by the number




Fig. 1. Chemical structure of hydroxytryptophan (Trp-OH), *N*-formylkynurenine (NFK), and kynurenine (Kyn)—products and intermediates of the tryptophan oxidation [17]. Alternative structures with hydroxyl substitution at the benzene ring are plausible for Trp-OH and cannot be distinguished based on the mass shift. The mass shifts observed by MS relative to Trp are noted beneath the formulas.

t1.1 **Table 1**
 Peptide masses defined in the inclusion list for LC–MS analysis.

t1.2	Peptide variant	Modification	[M + 2H] ²⁺	[M + 2H] ³⁺
t1.4	ILQWEADFLSHLK	Unmodified	800.4301	533.9558
t1.5	ILQW(OH)EADFLSHLK	Hydroxy-Trp	808.4272	539.2875
t1.6	ILQW(NFK)EADFLSHLK	NFK	816.4250	544.6191
t1.7	ILQW(Kyn)EADFLSHLK	Kyn	802.4276	535.2875
t1.8				
t1.9	YSIVVAATASEAAPLQYLAPFTGASIGEWFR	Unmodified	1643.8427	1096.2309
t1.10	YSIVVAATASEAAPLQYLAPFTGASIGEW(OH)FR	Hydroxy-Trp	1651.8402	1101.5626
t1.11	YSIVVAATASEAAPLQYLAPFTGASIGEW(NFK)FR	NFK	1659.8377	1106.8942
t1.12	YSIVVAATASEAAPLQYLAPFTGASIGEW(Kyn)FR	Kyn	1645.8402	1097.5626
t1.13				
t1.14	WNSGNDETK	Unmodified	533.7254	356.1527
t1.15	W(OH)NSGNDETK	Hydroxy-Trp	541.7229	361.4844
t1.16	W(NFK)NSGNDETK	NFK	527.7254	352.1527
t1.17	W(Kyn)NSGNDETK	Kyn	533.7254	356.1527

183 of spectral matches to a decoy database [33]. Acceptance criteria and
 184 filters were set to achieve a false positive rate of 5%.

185 2.6. Absolute peptide quantification

186 For quantitative analysis of the oxidative modification, single reac-
 187 tion monitoring (SRM) was applied on a triple quadrupole mass spec-
 188 trometer [34]. Custom peptides were synthesized for Trp and Kyn
 189 variants of the peptide I^{*}LQWEADFLSHLK by Thermo Scientific using
 190 isoleucine residue with stable isotopes to induce a mass shift of
 191 7 Da to the peptides. For analysis tryptic digests were spiked with
 192 peptide standards and analyzed on a Thermo TSQ Vantage coupled
 193 to the reversed phase chromatography described in Section 2.4. The
 194 SRM transitions used for the analysis are listed in Supplemental
 195 Table 1. For quantification of the Trp-OH and NFK variants the Trp refer-
 196 ence peptide was applied as standard, as only Trp and Kyn variants
 197 of the peptide could be obtained commercially. As judged by the sim-
 198 ilar ionization efficiencies observed for the Trp and Kyn variant, a sys-
 199 tematic error of less than 20% can be expected by this approximation.

200 3. Results

201 While indications for oxidative modification of tryptophan resi-
 202 dues in the ATP synthase in mitochondrial samples from *P. anserina*
 203 and other species have been described earlier [14], in this work
 204 focus was put on the detailed characterization, quantification, and lo-
 205 calization of these oxidative modifications within the protein com-
 206 plex. Crude mitochondria from juvenile (6 days) and senescent *P.*
 207 *anserina* cultures were isolated by differential centrifugation. For the
 208 investigation three independent cultures from each age stage were
 209 used. To determine the role of free iron as a factor for ROS generation,
 210 mitochondria were isolated from three independent *P. anserina* indi-
 211 viduals in the presence and absence of the strong iron chelating agent
 212 Desferal.

213 3.1. Separation of ATP synthase complexes

214 Fig. 2 displays the separation of solubilized mitochondrial mem-
 215 brane proteins by 2D-SDS/PAGE. BN-PAGE in the first dimension
 216 separates native monomers and dimers of the ATP synthase, while
 217 SDS-PAGE in the second dimension separates the protein subunits
 218 constituting these protein complexes. To estimate the apparent
 219 mass of the protein complexes in the blue-native gel digitonin-
 220 solubilized bovine heart mitochondria served as reference [14], as
 221 these are well characterized and are often used to estimate the
 222 mass of membrane protein complexes in blue-native gels [35, 36].
 223 In contrast to other species, e.g., rats [37], a separated F₁ subcomplex
 224 is hardly observed in the gels from *P. anserina* mitochondria. The gel
 225 system resolves the alpha subunits very well into two bands

226 originating from monomeric and dimeric ATP synthase complexes.
 227 The alpha subunit contains three tryptophan residues. Apart from
 228 the gamma subunit, which contains a single tryptophan, subunit
 229 alpha is the only subunit of the F₁ part containing tryptophan in its
 230 mature protein sequence.

231 3.2. Identification of tryptophan oxidation site

232 For localization and structural characterization of tryptophan oxi-
 233 dation sites in the primary structure of the alpha subunit, bands cor-
 234 responding to the dimeric and monomeric ATP synthase in the blue-
 235 native gel (Fig. 2: V₂, V₁) were digested with trypsin and the resulting
 236 tryptic peptides were applied to LC–MS/MS analysis. On average
 237 about 6500 MS/MS spectra could be assigned to peptides of ATP
 238 synthase subunits, yielding excellent sequence coverage of 30–80%
 239 for all proteins of the F₁ part. Although all subunits of the membrane
 240 integral F₀ part were also detected, the number of distinct peptides
 241 observed was significantly lower due to the reduced number of tryptic
 242 cleavage sites in these proteins [38]. For alpha and beta subunits
 243 sequence coverage of 50% and 72% respectively, was obtained with




Fig. 2. Monomers and dimers of ATP synthase from *P. anserina* mitochondria are efficiently resolved in 2D-BN/SDS-gels. In contrast to mitochondrial samples from other species, significant amounts of F₁ subcomplex are not observed in the gels.





Fig. 4. Relative proportion of different variants of the peptide ILQWEADFLSHLK quantified by the use of isotope-labeled reference peptides.

more than 2500 spectra assigned to peptides of each subunit. Fig. 3A displays a basepeak chromatogram of the analysis. All three Trp sites in the alpha subunit (Table 1) were sequenced by LC–MS/MS spectra from the corresponding tryptic peptides. Trp oxidation, however, could only be observed for Trp⁵⁰³ in the peptide ILQWEADFLSHLK. Species of this peptide with three intermediates of tryptophan oxidation—hydroxyl-tryptophan (Trp–OH), N-formylkynurenine (NFK), and kynurenine (Kyn) (Fig. 1)—were identified with p-values of $2 \cdot 10^{-6}$ and better. The assignment of these intermediates is unambiguous due to the high quality of the obtained MS/MS spectra (Fig. 3B–E). While the oxidation of the peptide could be demonstrated merely by the mass shift of the precursor and an incomplete ion series in the fragment spectra, the presence of y_9 and y_{10} ions unambiguously proves the oxidation of the Trp residue. The lower retention times as compared to the unmodified peptide (Trp: 68 min, Trp–OH: 51 min, NFK: 63 min, Kyn: 45 min) depict the reduced hydrophobicities of the oxidized Trp species.

Two other peptides of the alpha subunit containing Trp residues (Table 1) were observed, however, any corresponding peptides with oxidized Trp species could not be detected even when an inclusion list (Table 1) was applied for the LC–MS analysis.

3.3. Quantitation of tryptophan oxidation

For a quantitative analysis of the different oxidation intermediates of Trp⁵⁰³ observed in the LC–MS analysis isotope-labeled reference peptides were applied [34]. Fig. 4 depicts the relative proportion of the different oxidation intermediates of the peptide ILQWEADFLSHLK. Under all investigated conditions the NFK form was the most abundant oxidized species.

While the location of the oxidized tryptophan was well defined in all analyzed replicates, and also the amount of oxidized Trp was reproducible in technical replicates, as judged by peak areas of the respective peptides, a significant variation in the amount of oxidized Trp was observed between different biological replicates. In respect to the biological variation (for juvenile and senescent cultures), as well as monomeric and dimeric ATP synthase complexes, no statistically significant difference in the degree of oxidation was detected in the analysis in juvenile versus senescent cultures nor in monomeric versus dimeric ATP synthase. The degree of oxidation was, however, significantly reduced by more than 30%, when the strong iron

Fig. 3. A) Basepeak chromatogram of a tryptic digest from ATP synthase isolated from BN-gels analysed by LC–MS/MS. The four variants of the peptide ILQWEADFLSHLK are eluted with retention times of 68 min (Trp), 51 min (Trp–OH), 63 min (NFK), and 45 min (Kyn). Oxidized species from the two other tryptophan residues in the ATP synthase alpha subunit are not observed in the analysis. B–E) MS/MS spectra of the four variants of the peptide. The fragment ions y_9 and y_{10} representing the transition from EADFLSHLK to W^* EADFLSHLK allow the differentiation of oxidation intermediates.

283 chelator Desferal was present during the isolation of mitochondria
284 suggesting a potential involvement of iron or another transition
285 metal in the oxidation process.

286 4. Discussion

287 The ATP synthase is one of the most abundant proteins in the
288 inner mitochondrial membrane generating most of the approximat-
289 ely 70 kg ATP utilized in humans every day. It is devoid of redox-active
290 prosthetic groups and does not conduct any redox functions. Howev-
291 er, the regulation of the chloroplast ATP synthase by the redox state is
292 well known [39] and also for the mitochondrial enzyme indications
293 for a redox regulation have been shown recently [40]. Due to its spa-
294 tial proximity to the mitochondrial electron transport chain, the ATP
295 synthase is a primary target of ROS.

296 4.1. Selectivity of tryptophan oxidation

297 Using an LC–MS/MS approach the selective targeting of Trp⁵⁰³ of
298 the ATP synthase alpha subunit for oxidative modification can be un-
299 ambiguously displayed, while no oxidation can be observed for the
300 other Trp residues in the protein complex. Although oxidative stress
301 appears as an unstructured and random process, channeling of oxida-
302 tive damage to defined locations within protein complexes and pro-
303 tein subunits appears as a common characteristic of proteins frequently
304 exposed to ROS. This selectivity has been observed for mitochondrial
305 proteins—like the aconitase from bovine mitochondria displaying a pre-
306 dominant oxidation of Trp³⁷³ [15], as well as for proteins from other sys-
307 tems and organelles as the selective N-formylkynureninylation of Trp³⁶⁵
308 identified in subunit CP43 of the photosynthetic PSII complex under
309 high-light stress [20]. Recently, a targeting of oxidative modifications to
310 specific tryptophan residues was also recognized in pharmaceutical for-
311 mulations [41] and a detailed analysis could attribute selectivity to a
312 metal binding site in the proximity of the Trp residue.

313 4.2. Susceptibility of tryptophan towards oxidation

314 Although the oxidation of tryptophan is regularly observed in iso-
315 lated proteins, in contrast to cysteine and methionine the reactivity
316 towards superoxide anion radicals and hydrogen peroxide is quite
317 low. Taylor and coworkers observed the different susceptibility of
318 tryptophan and methionine towards oxidation, when analyzing pep-
319 tides from the 18 kDa subunit of complex I [17], and suggested differ-
320 ent oxidation mechanisms for the two amino acids. In pharmaceutical
321 products the low reactivity of H₂O₂ and peroxides with tryptophan
322 [42] is documented, while the reactivity of Trp with singlet oxygen

[43] and hydroxyl radicals released by the Fenton reaction is high
and is only surpassed by cysteine and methionine [44].

4.3. Localization within the ATP synthase structure

Fig. 5A displays a structural representation of the F₁ portion of the
ATP synthase. Due to the high homology with a sequence identity of
80%, based on the ATP synthase structure from *S. cerevisiae* [45] a
structure of subunit alpha from *P. anserina* was generated by homol-
ogy modeling using Swiss-Model [46–48]. In the resulting structure, it
is evident that all three tryptophans of subunit alpha are exposed to
the surface and potentially accessible from the surrounding. Trp⁵⁰³,
which is the sole target for oxidative modification, is located in the
peripheral region of the alpha subunit at a remote distance to catalyt-
ic and regulatory nucleotide binding sites or inter-subunit interfaces.
A feature distinguishing Trp⁵⁰³ from the other two Trp sites is the lo-
cation of three amino acid residues—Glu⁵⁰⁴, Asp⁵⁰⁶ and His⁵¹⁰ in its
proximity (Fig. 5B). These residues are able to constitute a binding
site for transition metal ion and induce an increased local concentra-
tion of these ions in the proximity of Trp⁵⁰³. While this sequence pat-
tern is well conserved in the subphylum Pezizomycotina and present
in 55 of 59 sequences available for subunit alpha from this subdivi-
sion in the NCBI database, it appears to be absent in unicellular
fungi in the subdivision Saccharomycotina and higher eukaryotes, in
which Trp⁵⁰³ is replaced by a phenylalanine in all known sequences.

4.4. Biological implications

As removal of damaged protein is a critical function for the main-
tenance of mitochondria [49] and an impairment of protein degrada-
tion and repair has been connected to age-associated accumulation of
oxidized proteins, for cellular homeostasis efficient means for the
perception of oxidative stress are required. While oxidized cysteine
and methionine can be efficiently reduced by repair mechanisms
[9], tryptophan oxidation products are irreversible and have the po-
tential to form markers visible for the mitochondrial quality control
system. While an effect of the oxidation of Trp⁵⁰³ on the enzymatic
activity of the ATP synthase is unlikely due to the distance to inter-
subunit contact sites and catalytical as well as regulatory nucleotide
binding sites, the disturbance of its structural integrity might be suf-
ficient to trigger recognition by the protein quality control system.
The LON protease, as an important component in the degradation of
oxidized proteins in the mitochondrial matrix [50, 51], is induced by
the presence of oxidized proteins both in activity and expression
level [34, 39]. Although not shown yet, the quality of the mitochon-
drial ATP synthase may be controlled by proteases like iAAA protease
or mAAA protease inserted in the inner mitochondrial membrane.




Fig. 5. A) Trp residue 503 (red) is surface exposed and located in the peripheral region of the alpha subunit with a distance of more than 30 Å from both the catalytic and regulatory nucleotide binding sites. The two other Trp residues (blue) are also exposed to the surface of the protein and are also accessible from the solvent. For the generation of a structural model of the alpha subunit of *P. anserina* a Swiss modeller [46–48] was used based on the structure published for *S. cerevisiae* [45] (PDB ID: 2XOK). B) The residues Glu⁵⁰⁴, Asp⁵⁰⁶ and His⁵¹⁰ located in the proximity of Trp⁵⁰³ at a distance of 5–9 Å are highlighted in a close-up of the structure. These residues are able to constitute a metal ion binding site. No such site can be defined for the two other Trp residues in the protein. For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.

With its reactivity towards hydroxyl radicals and singlet oxygen, tryptophan oxidation might mediate a sensory function for strong ROS or in connection with the putative metal binding site could be involved in the perception of metal catalyzed protein oxidation. Due to its abundance and its location in the inner mitochondrial membrane the ATP synthase might be able to perceive compounds with diffusion ranges limited by short half-lives [52] and mediate an up-regulation of the protein turnover before more sensible components are affected.

The channeling of oxidative damage to specific subunits and domains is a common feature observed in different enzymes evolved under the evolutionary pressure imposed by oxidative stress, like the D1 protein of the photosystem II as a prominent example [53] which accounts for up to 50% of protein synthesis [54, 55] in chloroplasts under high-light conditions. This focuses the effort for maintenance of protein complexes to the degradation and replacement of a single protein subunit [56] and, thereby transforms the random and untargeted process releasing ROS into a unidirectional signal that can be perceived by the quality control system.

Supplementary materials related to this article can be found online at doi:10.1016/j.bbabi.2011.11.006.

Acknowledgements

This work was funded by the German Federal Ministry for Education and Research (BMBF) through the *GerontoMitoSys* project (FKZ 0315584) to HDO, NAD and MR, and by the European Commission via a grant to NAD and HDO (acronym MiMage; LSHM-CT-2004-512020).

References

- [1] D. Harman, Aging—a theory based on free-radical and radiation-chemistry, *J. Gerontol.* 11 (1956) 298–300.
- [2] D. Harman, The biologic clock: the mitochondria? *J. Am. Geriatr. Soc.* 20 (1972) 145–147.
- [3] H.D. Osiewacz, Molecular analysis of aging processes in fungi, *Mutat. Res.* 237 (1990) 1–8.
- [4] E. Espagne, O. Lespinet, F. Malagnac, C. Da Silva, O. Jaillon, B.M. Porcel, A. Couloux, J.M. Aury, B. Segurens, J. Poulain, V. Anthouard, S. Grossetete, H. Khalili, E. Coppin, M. Dequard-Chablat, M. Picard, V. Contamine, S. Arnaise, A. Bourdais, V. Berteaux-Lecellier, D. Gautheret, R.P. de Vries, E. Battaglia, P.M. Coutinho, E.G. Danchin, B. Henrissat, R.E. Khoury, A. Sainsard-Chanet, A. Boivin, B. Pinan-Lucarre, C.H. Sellem, R. Debuchy, P. Wincker, J. Weissenbach, P. Silar, The genome sequence of the model ascomycete fungus *Podospora anserina*, *Genome Biol.* 9 (2008) R77.
- [5] H.D. Osiewacz, A. Hamann, DNA reorganization and biological aging. A review, *Biochemistry (Mosc)* 62 (1997) 1275–1284.
- [6] H.D. Osiewacz, Genetic regulation of aging, *J. Mol. Med.* 75 (1997) 715–727.
- [7] H.D. Osiewacz, E. Kimpel, Mitochondrial–nuclear interactions and lifespan control in fungi, *Exp. Gerontol.* 34 (1999) 901–909.
- [8] H.D. Osiewacz, Aging in fungi: role of mitochondria in *Podospora anserina*, *Mech. Ageing. Dev.* 123 (2002) 755–764.
- [9] S. Lorin, E. Dufour, A. Sainsard-Chanet, Mitochondrial metabolism and aging in the filamentous fungus *Podospora anserina*, *Biochim. Biophys. Acta* 1757 (2006) 604–610.
- [10] C.Q. Scheckhuber, H.D. Osiewacz, *Podospora anserina*: a model organism to study mechanisms of healthy ageing, *Mol. Genet. Genomics* 280 (2008) 365–374.
- [11] C. Borghouts, C.Q. Scheckhuber, A. Werner, H.D. Osiewacz, Respiration, copper availability and SOD activity in *P. anserina* strains with different lifespan, *Biogerontology* 3 (2002) 143–153.
- [12] R. Gredilla, J. Grief, H.D. Osiewacz, Mitochondrial free radical generation and lifespan control in the fungal aging model *Podospora anserina*, *Exp. Gerontol.* 41 (2006) 439–447.
- [13] S. Zintel, D. Schwitalla, K. Luce, A. Hamann, H.D. Osiewacz, Increasing mitochondrial superoxide dismutase abundance leads to impairments in protein quality control and ROS scavenging systems and to lifespan shortening, *Exp. Gerontol.* 45 (2010) 525–532.
- [14] K. Groebe, F. Krause, B. Kunstmann, H. Unterluggauer, N.H. Reifschneider, C.Q. Scheckhuber, C. Sastri, W. Stegmann, W. Wozny, G.P. Schwall, S. Poznanovic, N.A. Dencher, P. Jansen-Durr, H.D. Osiewacz, A. Schratzenholz, Differential proteomic profiling of mitochondria from *Podospora anserina*, rat and human reveals distinct patterns of age-related oxidative changes, *Exp. Gerontol.* 42 (2007) 887–898.
- [15] C. Hunzinger, W. Wozny, G.P. Schwall, S. Poznanovic, W. Stegmann, H. Zengerling, R. Schoepf, K. Groebe, M.A. Cahill, H.D. Osiewacz, N. Jagemann, M. Bloch, N.A. Dencher, F. Krause, A. Schratzenholz, Comparative profiling of the mammalian

- mitochondrial proteome: multiple aconitase-2 isoforms including N-formylkynurenine modifications as part of a protein biomarker signature for reactive oxidative species, *J. Proteome Res.* 5 (2006) 625–633.
- [16] B.S. Berlett, E.R. Stadtman, Protein oxidation in aging, disease, and oxidative stress, *J. Biol. Chem.* 272 (1997) 20313–20316.
- [17] S.W. Taylor, E. Fahy, J. Murray, R.A. Capaldi, S.S. Ghosh, Oxidative post-translational modification of tryptophan residues in cardiac mitochondrial proteins, *J. Biol. Chem.* 278 (2003) 19587–19590.
- [18] M. Fedorova, T. Todorovsky, N. Kuleva, R. Hoffmann, Quantitative evaluation of tryptophan oxidation in actin and troponin I from skeletal muscles using a rat model of acute oxidative stress, *Proteomics* 10 (2010) 2692–2700.
- [19] E.L. Finley, J. Dillon, R.K. Crouch, K.L. Schey, Identification of tryptophan oxidation products in bovine alpha-crystallin, *Protein Sci.* 7 (1998) 2391–2397.
- [20] T.M. Dreaden, J. Chen, S. Rexroth, B.A. Barry, N-formylkynurenine as a marker of high light stress in photosynthesis, *J. Biol. Chem.* 286 (2011) 22632–22641.
- [21] S. Rinalducci, N. Campostrini, P. Antonoli, P.G. Righetti, P. Roepstorff, L. Zolla, Formation of truncated proteins and high-molecular-mass aggregates upon soft illumination of photosynthetic proteins, *J. Proteome Res.* 4 (2005) 2327–2337.
- [22] D. Brust, A. Hamann, H.D. Osiewacz, Deletion of PaAif2 and PaAmid2, two genes encoding mitochondrial AIF-like oxidoreductases of *Podospora anserina*, leads to increased stress tolerance and lifespan extension, *Curr. Genet.* 56 (2010) 225–235.
- [23] C. Borghouts, S. Kerschner, H.D. Osiewacz, Copper-dependence of mitochondrial DNA rearrangements in *Podospora anserina*, *Curr. Genet.* 37 (2000) 268–275.
- [24] H. Schägger, G. von Jagow, Blue native electrophoresis for isolation of membrane protein complexes in enzymatically active form, *Anal. Biochem.* 199 (1991) 223–231.
- [25] H.J. Schwassmann, S. Rexroth, H. Seelert, N.A. Dencher, Metabolism controls dimerization of the chloroplast FoF1 ATP synthase in *Chlamydomonas reinhardtii*, *FEBS Lett.* 581 (2007) 1391–1396.
- [26] H. Schägger, W.A. Cramer, G. von Jagow, Analysis of molecular masses and oligomeric states of protein complexes by blue native electrophoresis and isolation of membrane protein complexes by two-dimensional native electrophoresis, *Anal. Biochem.* 217 (1994) 220–230.
- [27] T. Rabilloud, G. Carpentier, P. Tarroux, Improvement and simplification of low-background silver staining of proteins by using sodium dithionite, *Electrophoresis* 9 (1988) 288–291.
- [28] H. Blum, H. Beier, H.J. Gross, Improved silver staining of plant-proteins, RNA and DNA in polyacrylamide gels, *Electrophoresis* 8 (1987) 93–99.
- [29] F. Gharahdaghi, C.R. Weinberg, D.A. Meagher, B.S. Imai, S.M. Mische, Mass spectrometric identification of proteins from silver-stained polyacrylamide gel: a method for the removal of silver ions to enhance sensitivity, *Electrophoresis* 20 (1999) 601–605.
- [30] S. Rexroth, J.M. Meyer Zu Tittingdorf, F. Krause, N.A. Dencher, H. Seelert, Thylakoid membrane at altered metabolic state: challenging the forgotten realms of the proteome, *Electrophoresis* 24 (2003) 2814–2823.
- [31] J. Oetjen, S. Rexroth, B. Reinhold-Hurek, Mass spectrometric characterization of the covalent modification of the nitrogenase Fe-protein in *Azoarcus sp* BH72, *FEBS J.* 276 (2009) 3618–3627.
- [32] J.K. Eng, A.L. McCormack, J.R. Yates III, An approach to correlate tandem mass-spectral data of peptides with amino-acid-sequences in a protein database, *J. Am. Soc. Mass Spectrom.* 5 (1994) 976–989.
- [33] L. Kall, J.D. Storey, M.J. MacCoss, W.S. Noble, Assigning significance to peptides identified by tandem mass spectrometry using decoy databases, *J. Proteome Res.* 7 (2008) 29–34.
- [34] S.A. Gerber, J. Rush, O. Stemman, M.W. Kirschner, S.P. Gygi, Absolute quantification of proteins and phosphoproteins from cell lysates by tandem MS, *Proc. Natl. Acad. Sci. USA* 100 (2003) 6940–6945.
- [35] L. Jänsch, V. Kruff, U.K. Schmitz, H.P. Braun, Unique composition of the preprotein translocase of the outer mitochondrial membrane from plants, *J. Biol. Chem.* 273 (1998) 17251–17257.
- [36] S. Rexroth, J.M. Meyer Zu, H.J. Tittingdorf, F. Schwassmann, H. Krause, N.A. Seelert, Dencher, Dimeric H+–ATP synthase in the chloroplast of *Chlamydomonas reinhardtii*, *Biochim. Biophys. Acta* 1658 (2004) 202–211.
- [37] M. Frenzel, H. Rommelspacher, M.D. Sugawa, N.A. Dencher, Ageing alters the supramolecular architecture of OxPhos complexes in rat brain cortex, *Exp. Gerontol.* 45 (2010) 563–572.
- [38] A. Poetsch, S. Rexroth, J. Heberle, T.A. Link, N.A. Dencher, H. Seelert, Characterisation of subunit III and its oligomer from spinach chloroplast ATP synthase, *Biochim. Biophys. Acta* 1618 (2003) 59–66.
- [39] R.E. McCarty, ATP synthase of chloroplast thylakoid membranes: a more in depth characterization of its ATPase activity, *J. Bioenerg. Biomembr.* 37 (2005) 289–297.
- [40] S.B. Wang, D.B. Foster, J. Rucker, B. O'Rourke, D.A. Kass, J.E. Van Eyk, Redox regulation of mitochondrial ATP synthase: implications for cardiac resynchronization therapy, *Circ. Res.* (2011).
- [41] X.M. Lam, W.G. Lai, E.K. Chan, V. Ling, C.C. Hsu, Site-specific tryptophan oxidation induced by autocatalytic reaction of polysorbate 20 in protein formulation, *Pharm. Res.* (2011).
- [42] J.A. Ji, B. Zhang, W. Cheng, Y.J. Wang, Methionine, tryptophan, and histidine oxidation in a model protein, PTH: mechanisms and stabilization, *J. Pharm. Sci.* 98 (2009) 4485–4500.
- [43] M.J. Davies, Reactive species formed on proteins exposed to singlet oxygen, *Photochem. Photobiol. Sci.* 3 (2004) 17–25.
- [44] G. Xu, M.R. Chance, Radiolytic modification and reactivity of amino acid residues serving as structural probes for protein footprinting, *Anal. Chem.* 77 (2005) 4549–4555.

- 524 [45] D. Stock, A.G. Leslie, J.E. Walker, Molecular architecture of the rotary motor in ATP
525 synthase, *Science (New York, N.Y.)* 286 (1999) 1700–1705. 537
- 526 [46] K. Arnold, L. Bordoli, J. Kopp, T. Schwede, The SWISS-MODEL workspace: a web-
527 based environment for protein structure homology modelling, *Bioinformatics*
528 22 (2006) 195–201. 538
- 529 [47] F. Kiefer, K. Arnold, M. Kunzli, L. Bordoli, T. Schwede, The SWISS-MODEL reposi-
530 tory and associated resources, *Nucleic Acids Res.* 37 (2009) D387–D392. 539
- 531 [48] M.C. Peitsch, Protein modeling by e-mail, *Nat. Biotechnol.* 13 (1995) 658–660. 540
- 532 [49] I. Petropoulos, B. Friguet, Maintenance of proteins and aging: the role of oxidized
533 protein repair, *Free. Radic. Res.* 40 (2006) 1269–1276. 541
- 534 [50] D.A. Bota, K.J. Davies, Lon protease preferentially degrades oxidized mitochon-
535 drial aconitase by an ATP-stimulated mechanism, *Nat. Cell Biol.* 4 (2002)
536 674–680. 542
- [51] K. Luce, H.D. Osiewacz, Increasing organismal healthspan by enhancing mito-
chondrial protein quality control, *Nat. Cell Biol.* 11 (2009) 852–858. 543
- [52] W.A. Pryor, Oxy-radicals and related species: their formation, lifetimes, and reac-
tions, *Annu. Rev. Physiol.* 48 (1986) 657–667. 544
- [53] I. Ohad, D.J. Kyle, C.J. Arntzen, Membrane-protein damage and repair—removal
and replacement of inactivated 32-kilodalton polypeptides in chloroplast mem-
branes, *J. Cell Biol.* 99 (1984) 481–485. 545
- [54] J.M. Anderson, W.S. Chow, Structural and functional dynamics of plant photosys-
tem II, *Philos. Trans. R. Soc. Lond. Ser. B-Biol. Sci.* 357 (2002) 1421–1430. 546
- [55] J. Barber, B. Andersson, Too much of a good thing—light can be bad for photosyn-
thesis, *Trends Biochem. Sci.* 17 (1992) 61–66. 547
- [56] A.W. Rutherford, P. Faller, Photosystem II: evolutionary perspectives, *Philos.*
548 *Trans. R. Soc. Lond. Ser. B-Biol. Sci.* 358 (2003) 245–253. 549
- 550

551

UNCORRECTED PROOF